

newASTROGAM – The New MeV to GeV Gamma-ray Observatory

D. Berge,^{a,b,*} M. N. Mazziotta,^c M. Tavani,^d V. Tatischeff^e and U. Oberlack^f on behalf of the newASTROGAM proposal team

^a*Deutsches Elektronen-Synchrotron DESY, Platanenallee 6, D-15738 Zeuthen, Germany*

^b*Humboldt-University of Berlin, Institute of Physics, Newtonstraße 15, D-12489 Berlin, Germany*

^c*Istituto Nazionale di Fisica Nucleare, Sezione di Bari, via Orabona 4, I-70126 Bari, Italy*

^d*INAF-IAPS Roma, via del Fosso del Cavaliere 100, I-00133 Roma, Italy*

^e*Université Paris-Saclay, CNRS/IN2P3, IJCLab, F-91405 Orsay, France*

^f*Institut für Physik & Exzellenzcluster PRISMA+, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany*

E-mail: david.berge@desy.de

newASTROGAM is a breakthrough mission concept for the study of the non-thermal Universe from space with γ -rays in the energy range from 15 keV to 3 GeV. It is based on advanced space-proven detector technologies, which will achieve unprecedented sensitivity, angular and energy resolution combined with polarimetric capability. Since the MeV γ -ray energy range is the most under-explored electromagnetic window to the Universe, a mission in this energy range can for the first time sensitively address fundamental astrophysics questions connected to the physics of compact objects and merger events, jets and their environments, supernovae and the origin of the elements, potentially constrain the nature of dark matter and many more science objectives. The mission will detect and follow-up many of the key sources of multi-messenger astronomy in the 2040s.

newASTROGAM provides an unprecedentedly broad energy coverage from keV to GeV energies. The payload concept consists of a Silicon tracker combined with a crystal calorimeter. Both detectors are surrounded by an anti-coincidence detector to reject charged cosmic rays. In addition, a thin X-ray coded mask provides very good imaging capabilities. Such a mission can uniquely detect γ -rays via the photoelectric effect, Compton scattering and electron-positron pair production. newASTROGAM is proposed to the ESA call for medium-class mission ideas (M8).

39th International Cosmic Ray Conference (ICRC2025)
15–24 July 2025
Geneva, Switzerland



*Speaker

1. Introduction

The γ -ray sky at MeV and GeV energies provides unique and crucial information on the most energetic and intriguing phenomena of our Universe. Photons in this range provide fundamental insights into the physics of nucleosynthesis, accretion, particle acceleration, strong magnetic fields, and extreme-gravity environments [1]. Observing the sky at these energies with a large field of view (FoV) and unprecedentedly high sensitivity is of paramount importance in the era of multi-messenger astronomy, as clearly underlined in the [Voyage 2050](#) program and [Astro2020 Decadal Survey](#) report.

*newASTROGAM*¹ is designed as a unique multi-purpose observatory with scanning and pointing capability, enabling imaging, polarimetry, line and continuum spectroscopy, sub-millisecond timing, and fast transient detection in the MeV-GeV energy domain. It will be the first instrument with adequate sensitivity in the MeV range and simultaneous GeV coverage, thus ensuring the broad-band information needed to unravel many fundamental physical processes. *newASTROGAM* is *new* because a hard-X-ray imager with arcminute source localization capability has been added to the original design. This addition broadens the purpose of the mission, which will not only complement data from, but also provide trigger information to, prominent facilities such as SKA, ALMA, Roman Space Telescope, ELT, newAthena, CTAO, SWGO, LISA, the Einstein Telescope, IceCube-Gen2, KM3NeT.

While the mission will contribute to many science topics of interest for a broad and diverse community, *newASTROGAM* promises major breakthroughs in the following areas:

- physics of extreme cosmic accelerators from within the Galaxy to Cosmic Dawn;
- nucleosynthesis, supernova explosions, chemical evolution of the Interstellar Medium (ISM);
- Cosmic Ray (CR) sources and feedback;
- the extreme Universe in the era of multi-messenger astronomy;
- fundamental physics.

2. Science

With its large FoV ($\sim 1/3$ of the sky), all-sky scanning mode, and arcmin localization capability, *newASTROGAM* will study the activity of thousands of cosmic accelerators such as pulsars, stellar and supermassive black holes (SMBHs), γ -ray bursts (GRBs), magnetars, supernovae (SNe), and kilonovae. With its unprecedented sensitivity (see Fig. 1), it will provide the deepest all-sky mapping of the MeV emission produced by the radioactive decay of newly synthesized nuclei, CR interactions, $e^\pm$ pairs, and decisively search the MeV-GeV sky for dark-matter (DM) signatures. Finally, it will help identify electromagnetic counterparts to gravitational-wave (GW) and high-energy neutrino sources, expanding the pioneering role of the *Fermi* satellite.

Insights into Extreme Acceleration Processes

MeV-GeV variability studies of relativistic jets can be conducted with *newASTROGAM* on timescales of seconds to hours for GRBs and from minutes to years for AGN. For AGN jets they will provide

¹Mission homepage: <https://www.new-astrogam.eu>




Figure 1: Point-source (extra-galactic) continuum sensitivity of different X- and γ -ray instruments.

crucial information on the size and location of the acceleration site, and on the Lorentz factor and any precession of the jet. The detailed shape of the emission spectrum around the MeV peak places important constraints on acceleration and radiation mechanisms, and maximum particle energy [2]. *newASTROGAM* data will also constrain the lepton/baryon content and magnetization of the jets, whose knowledge is fundamental to assessing their energetics, the microphysics of particle acceleration, and the impact on their surroundings. Analogous constraints on the sites and mechanisms of particle acceleration will be placed on black-hole (BH) and neutron-star (NS) X-ray binaries, recently probed by INTEGRAL and *Fermi* as high-energy emitters, with the MeV band providing the missing link [3, 4].

γ -ray polarimetry [5] is a young field with a large discovery potential. Polarisation changes between X-rays and γ -rays can probe the location and evolution of relativistic particle outflows. *newASTROGAM* will enable polarimetric measurements for tens of Galactic sources with sensitivity to variations in time. For accreting BHs, measurements of the polarisation and the MeV cutoff energy can firmly establish the nature of the observed MeV tail, disentangling its origin between the jet and the disk corona.

Current observations of TeV-to-PeV neutrinos suggest the existence of both a Galactic and an extragalactic population of hadronic PeVatrons (PeV ion accelerators) mostly hidden to GeV-TeV γ -ray telescopes due to strong γ -ray absorption in the source. A handful of temporal associations between neutrinos and γ -ray flares suggest AGN jets and Galactic γ -ray binaries as promising PeVatrons [6]. Some steady neutrino sources [7] are claimed to be associated with AGN coronae. The accompanying cascade radiation would preferentially appear in the MeV band [8], making *newASTROGAM*'s role decisive to decipher the processes of PeV particle acceleration, both in Seyfert galaxies [9] and in blazars [10].

newASTROGAM will shed light on BH activity in the early Universe. The most luminous AGN,

powered by accretion onto a SMBH of about $10^9 M_\odot$, are most abundant at redshifts $z \gtrsim 3$ [11], before the cosmic noon in star-formation history. The jet emission peaks in the MeV-GeV band, and largely dominates the radiative output when the jet points in our direction (blazar). *newASTROGAM* can characterise the jet activity of hundreds of objects, both in quiescence and during flares. MeV blazars can be detected up to redshift $z \sim 6 - 7$ [11], though most of the detected sources will have redshifts $z < 4$. The opportunity is unique to shed light on the physics of this source class, the cosmological evolution of SMBHs, the environment of their growth, and the extent to which AGN jets influence galaxy evolution [12]. This topic is also identified as a Voyage 2050 priority.

Explosive Nucleosynthesis and Chemical Evolution of Galaxies




Figure 2: Improvement over INTEGRAL (top, [13]) for a type-Ia SN. Lines from the ^{56}Ni decay chain will clearly be resolved (bottom), yielding the radioactive and total ejecta masses, a characteristic of the progenitors.

Interstellar enrichment in heavy nuclei is driven by asymptotic-giant-branch stars, core-collapse supernovae (SNe), white-dwarf explosions, and NS mergers (kilonovae), but their relative contributions are not established at all [14]. *newASTROGAM* can detect important nuclear decay lines with unprecedented sensitivity and, for nominal mission lifetimes, improve over the upcoming COSI mission [15] by factors 2.5 to 30, depending on the line.

The large FoV will help detecting γ -rays, the earliest photons to escape explosion sites, before the optical peak, for optimal constraints on the nature and mass of the progenitor and on the explosion itself [16]. The detection of type-Ia SNe out to a distance of 30 Mpc will enable precise

measurements of the total mass of $^{56}\text{Ni}/^{56}\text{Co}$ in the ejecta (see Fig. 2), probing the progenitors and also testing the calibration of the cosmological distance determination with SNe Ia [17]. γ -rays from the $^{44}\text{Ti}/^{44}\text{Sc}$ decay chain can be detected in most of the young Galactic supernova remnants and in SN1987A, providing information on the degree of asymmetry and clumpiness of the ejecta of core-collapse SNe [1].

newASTROGAM can detect line emission from kilonovae up to 12 Mpc away and measure the mass of nuclei synthesised by rapid neutron capture. In the Milky Way, it can uncover the remnant of a kilonova up to 10–100 kyr old through long-lived radioisotopes such as ^{126}Sn and their fission products [18–20]. *newASTROGAM* will map the positron annihilation line at 511 keV with unprecedented quality and may even obtain the first point-source detection. It can detect the 2.2-MeV neutron-capture line from accreting neutron stars. It can also map the long-lived ^{26}Al and ^{60}Fe radioisotopes, shedding new light on stellar nucleosynthesis and on the subsequent mixing of the high-metallicity ejecta into the ISM [21]. Finally, it may provide the first unambiguous estimate of the ^{22}Na and ^7Be (see ref. [22] for INTEGRAL hints of the latter) yield in a nova, and thus gauge the nova contribution to the Galactic enrichment in ^7Li .

Cosmic Ray Sources and Feedback on Galaxy Evolution

newASTROGAM data will be crucial to resolve important CR puzzles. Observations of CR accelerators below 1 GeV will be decisive for disentangling the emission of accelerated ions and electrons, and that of freshly accelerated versus re-accelerated CRs, and will allow studying CR feedback on the sources. *newASTROGAM* will provide the first precise measurement of the “Pion Bump” below 500 MeV, a direct probe of hadronic emission [23]. MeV-band spectra of CR sources will probe the initial acceleration of CR electrons and hence the connection between CRs and quasi-thermal plasma [24], while nuclear de-excitation lines will provide otherwise inaccessible information on CR ions below 200 MeV/nucleon.

The impact of CRs is a highly uncertain feedback factor in galaxy evolution. The pressure gradient of GeV CRs can drive galactic outflows, whereas low-energy (< 100 MeV) CRs are the main source of ionisation and heating in dark clouds, with direct impact on star formation. *newASTROGAM* observations will advance our poor understanding of the propagation of these two important CR populations, probing their spectra at different locations in the disk, near their sources, in the “quiet” ISM and in the highly turbulent medium of starburst regions, where re-acceleration can occur [25, 26]. Mapping their large-scale distribution will measure beyond the Galactic disk the imprint of star-forming regions [27].

In the Galactic center region, *newASTROGAM* will improve the spatial and spectral characterisation of the puzzling γ -ray excess, to elucidate its debated origin between DM annihilation, an excess of CRs, or a population of unresolved millisecond pulsars [28]. On larger scales, the angular resolution and energy band of the instrument are ideal to study the morphology and ion/lepton composition of the Fermi bubbles, and help assess their origin as a CR-driven wind or as jets powered by the central BH, Sgr A* [29, 30].

Deciphering the Energetic Transient Sky

The sensitivity, FoV, and spectral range of *newASTROGAM* will allow the MeV/GeV-band detection of hundreds of long (see Fig. 3) and short GRBs. The latter, counterparts to binary NS mergers, will




Figure 3: X- and γ -ray afterglow spectrum of a long GRB [31]. *newASTROGAM* probes the poorly covered transition region, its sensitivity is shown in red here for 600 s exposure.

be detected at a rate of several tens per year, ensuring synergy with GW detectors. For the former, the mission will not only see the bright prompt emission, but can uniquely follow the afterglow and determine the radiation process and the spectrum of the radiating particles.

The X-ray Coded Mask will yield arcmin localisation, guiding multi-wavelength follow-up observations of the source environment and of the development of the kilonova as softer light emerges hours and days later. Moreover, *newASTROGAM* will measure intermediate X- to γ -rays in a special burst event mode (*calorimeter count rates*) and provide polarimetry for the brightest events, key to identifying the radiation process and constrain the magnetic and kinetic profile of the jet [32]. Time-domain surveys conducted by SKA in the radio, Roman in the infrared, and Rubin in the optical, will revolutionise our knowledge of the transient sky: *newASTROGAM* will add the fundamental MeV-GeV band to this revolution, covering many source classes beyond the well-studied micro-quasars, AGN and GRB transients, including magnetars [33] and Fast Blue Optical Transients [34]. All detected transient signals will be rapidly disseminated with arcminute localization in hard X-rays.

Fundamental and New Physics

newASTROGAM will explore fundamental physics. It will access the MeV to GeV mass scale of DM, where direct searches are insensitive and collider searches are highly model dependent, and offer multiple angles for detections [35, 36]: its angular resolution allows DM signals from

subhalos and the inner Galaxy to be better identified than with *Fermi*, it offers unprecedented details of DM models explaining the 511 keV line [37, 38], and can differentiate pulsar and DM interpretations of the Galactic-center excess [35]. Weakly Interacting Sub-eV Particles, e.g. axions and axion-like particles (ALP) [39], would produce spectral signatures in the MeV band [40] or could be probed by ALP-induced GRBs from extragalactic SNe [41, 42]. Polarimetry of bright GRBs will address fundamental questions related to vacuum birefringence and Lorentz-invariance violation [43]. Finally, *newASTROGAM* will be highly sensitive to photon emission from primordial BHs in the mass range 10^{15-17} g, which peaks at 1-30 MeV [44, 45].

3. *newASTROGAM* Mission Configuration

To achieve a breakthrough sensitivity, *newASTROGAM* will be launched in an equatorial low-Earth orbit of inclination $< 2.5^\circ$, eccentricity < 0.01 , and altitude 550 – 600 km. Such an orbit is only marginally affected by the South Atlantic Anomaly therefore minimising instrumental particle backgrounds, ideal for high-energy observations [46]. Alerts for new transient sources detected onboard with arcminute accuracy will be instantly transmitted to the ground. The mission will mostly do survey observations covering with its large FoV the entire sky in one day. In addition, occasional target-of-opportunity observations can be performed by a nearly inertial pointing to focus on a particular region. The nominal mission duration is planned to be three years.

The payload consists of a single instrument operating over five orders of magnitude in photon energy (15 keV–3 GeV), detecting photons via the photoelectric effect, Compton scattering, and $e^\pm$ pair production. To veto cosmic rays, the outermost Anti-Coincidence (AC) system is composed of segmented panels of plastic scintillators covering the top and the lateral sides of the roughly quadratic payload (≈ 120 cm across). The AC vetoes more than 99.99% of the penetrating charged particles and is based on heritage of the *AGILE* and *Fermi* missions. Within the AC and 14 cm before the first Silicon Tracker layer is a thin Coded Mask to image X-rays in the ~ 15 –40 keV energy range with an angular resolution of $\sim 15'$, providing a localisation capability of a few arcminutes for bright sources. The mask consists of a tungsten sheet with a pseudo-random pattern (open fraction 50%). The mask mechanical support is also provided by an X-ray collimator of carbon fibre walls. The FoV of the X-ray coded mask monitor is 4.06 sr (half coded).

The instrument's heart is a Silicon microstrip Tracker. Such strips are already employed for γ -ray detection in space (e.g. for *DAMPE*). The Tracker measures hard X-rays that have passed the Coded Mask and γ -rays that undergo Compton scattering or pair conversion. The baseline design has double-sided detectors (DSSDs) read out by low-noise and low-power electronics with self-triggering capability. The Tracker is supported by a very light carbon fibre mechanical structure. 9 DSSDs are wire bonded strip to strip to form a 2-D ladder. Alternatively, CMOS pixels [47, 48] will be assessed and compared to the baseline DSSDs during the study phase.

The Calorimeter is a pixelated detector made of a high- Z scintillation material (such as CsI(Tl)) for efficient absorption of γ -rays and $e^\pm$ pairs. It consists of an array of parallelepiped crystal bars read out by silicon drift detectors or silicon photomultipliers. The depth of interaction in each crystal is measured from the difference of recorded scintillation signals at both ends. The Calorimeter architecture is based on the heritage of instruments like for example *INTEGRAL*/PICsIT or *POLAR-2*.

References

- [1] A. de Angelis, V. Tatischeff, I.A. Grenier, J. McEnery, M. Mallamaci, M. Tavani et al., *Science with e-ASTROGAM. A space mission for MeV-GeV gamma-ray astrophysics*, *Journal of High Energy Astrophysics* **19** (2018) 1 [1711.01265].
- [2] F. Tavecchio, L. Nava, A. Sciacaluga and P. Coppi, *Probing the low-energy particle content of blazar jets through MeV observations*, *A&A* **694** (2025) L3 [2412.09089].
- [3] A. Papitto and D. de Martino, *Transitional Millisecond Pulsars*, in *Astrophysics and Space Science Library*, S. Bhattacharyya, A. Papitto and D. Bhattacharya, eds., vol. 465 of *Astrophysics and Space Science Library*, pp. 157–200, Jan., 2022, DOI [2010.09060].
- [4] S.E. Motta, J. Rodriguez, E. Jourdain, M. Del Santo, G. Belanger, F. Cangemi et al., *The INTEGRAL view on black hole X-ray binaries*, *New A Rev.* **93** (2021) 101618 [2105.05547].
- [5] D. Götz, C. Gouiffès, J. Rodriguez, P. Laurent, E. Jourdain, J.-P. Roques et al., *INTEGRAL results on gamma-ray bursts and polarization of hard X-ray sources*, *New A Rev.* **87** (2019) 101537.
- [6] A.M. Bykov, A.E. Petrov, M.E. Kalyashova and S.V. Troitsky, *PeV Photon and Neutrino Flares from Galactic Gamma-Ray Binaries*, *ApJ* **921** (2021) L10 [2110.11189].
- [7] IceCube Collaboration, R. Abbasi, M. Ackermann, J. Adams, J.A. Aguilar, M. Ahlers et al., *Evidence for neutrino emission from the nearby active galaxy NGC 1068*, *Science* **378** (2022) 538 [2211.09972].
- [8] K. Murase, *Hidden Hearts of Neutrino Active Galaxies*, *ApJ* **941** (2022) L17 [2211.04460].
- [9] K. Murase, C.M. Karwin, S.S. Kimura, M. Ajello and S. Buson, *Sub-GeV Gamma Rays from Nearby Seyfert Galaxies and Implications for Coronal Neutrino Emission*, *ApJ* **961** (2024) L34 [2312.16089].
- [10] G. Ficht de Clairfontaine, S. Buson, L. Pfeiffer, S. Marchesi, A. Azzollini, V. Baghmanyany et al., *Hadronic Processes at Work in 5BZB J0630-2406*, *ApJ* **958** (2023) L2 [2310.03698].
- [11] L. Marcotulli, M. Ajello, C.M. Urry, V.S. Paliya, M. Koss, K. Oh et al., *BASS. XXXIII. Swift-BAT Blazars and Their Jets through Cosmic Time*, *ApJ* **940** (2022) 77 [2209.09929].
- [12] T. Sbarrato, *Big and Young Supermassive Black Holes in the Early Universe*, *Galaxies* **9** (2021) 23 [2107.09940].
- [13] E. Churazov, R. Sunyaev, J. Isern, I. Bikmaev, E. Bravo, N. Chugai et al., *Gamma-rays from Type Ia Supernova SN2014J*, *ApJ* **812** (2015) 62 [1502.00255].
- [14] J.A. Johnson, B.D. Fields and T.A. Thompson, *The origin of the elements: a century of progress*, *Philosophical Transactions of the Royal Society of London Series A* **378** (2020) 20190301.

- [15] J. Tomsick, S. Boggs, A. Zoglauer, D.H. Hartmann, M. Ajello, E. Burns et al., *The Compton Spectrometer and Imager*, in *38th International Cosmic Ray Conference*, p. 745, Sept., 2024, DOI [2308.12362].
- [16] J. Isern, M. Hernanz, E. Bravo, S. Grebenev, P. Jean, M. Renaud et al., *Synthesis of radioactive elements in novae and supernovae and their use as a diagnostic tool*, *New A Rev.* **92** (2021) 101606 [2101.02738].
- [17] M.M. Phillips, *The Absolute Magnitudes of Type IA Supernovae*, *ApJ* **413** (1993) L105.
- [18] M.-R. Wu, P. Banerjee, B.D. Metzger, G. Martínez-Pinedo, T. Aramaki, E. Burns et al., *Finding the Remnants of the Milky Way’s Last Neutron Star Mergers*, *ApJ* **880** (2019) 23 [1905.03793].
- [19] O. Korobkin, A.M. Hungerford, C.L. Fryer, M.R. Mumpower, G.W. Misch, T.M. Sprouse et al., *Gamma Rays from Kilonova: A Potential Probe of r-process Nucleosynthesis*, *ApJ* **889** (2020) 168 [1905.05089].
- [20] X. Wang, N3AS Collaboration, N. Vassh, FIRE Collaboration, T. Sprouse, M. Mumpower et al., *MeV Gamma Rays from Fission: A Distinct Signature of Actinide Production in Neutron Star Mergers*, *ApJ* **903** (2020) L3 [2008.03335].
- [21] R. Diehl, M. Lugaro, A. Heger, A. Sieverding, X. Tang, K.A. Li et al., *The radioactive nuclei ^{26}Al and ^{60}Fe in the Cosmos and in the solar system*, *PASA* **38** (2021) e062 [2109.08558].
- [22] L. Izzo, T. Siegert, P. Jean, P. Molaro, P. Bonifacio, M. Della Valle et al., *Possible evidence for the 478 keV emission line from ^7Be decay during the outburst phases of V1369 Cen*, *A&A* **698** (2025) A291 [2504.20866].
- [23] J. Liu, B. Liu and R. Yang, *Precise measurement of pion-bump structure using future MeV gamma-ray detectors*, *Phys. Rev. D* **110** (2024) 063026 [2406.03691].
- [24] T. Amano, Y. Matsumoto, A. Bohdan, O. Kobzar, S. Matsukiyo, M. Oka et al., *Nonthermal electron acceleration at collisionless quasi-perpendicular shocks*, *Reviews of Modern Plasma Physics* **6** (2022) 29 [2209.03521].
- [25] M. Ackermann, M. Ajello, A. Allafort, L. Baldini, J. Ballet, G. Barbiellini et al., *A Cocoon of Freshly Accelerated Cosmic Rays Detected by Fermi in the Cygnus Superbubble*, *Science* **334** (2011) 1103.
- [26] A.M. Bykov, A. Marcowith, E. Amato, M.E. Kalyashova, J.M.D. Kruijssen and E. Waxman, *High-Energy Particles and Radiation in Star-Forming Regions*, *Space Sci. Rev.* **216** (2020) 42 [2003.11534].
- [27] I.A. Grenier, J.H. Black and A.W. Strong, *The Nine Lives of Cosmic Rays in Galaxies*, *ARA&A* **53** (2015) 199.
- [28] M. Di Mauro, *Characteristics of the Galactic Center excess measured with 11 years of Fermi -LAT data*, *Phys. Rev. D* **103** (2021) 063029 [2101.04694].

- [29] M. Ackermann, A. Albert, W.B. Atwood, L. Baldini, J. Ballet, G. Barbiellini et al., *The Spectrum and Morphology of the Fermi Bubbles*, [ApJ **793** \(2014\) 64 \[1407.7905\]](#).
- [30] M. Negro, H. Fleischhack, A. Zoglauer, S. Digel and M. Ajello, *Unveiling the Origin of the Fermi Bubbles with MeV Photon Telescopes*, [ApJ **927** \(2022\) 225](#).
- [31] MAGIC Collaboration, V.A. Acciari, S. Ansoldi, L.A. Antonelli, A.A. Engels, D. Baack et al., *Observation of inverse Compton emission from a long γ -ray burst*, [Nature **575** \(2019\) 459 \[2006.07251\]](#).
- [32] E. Burns, *Neutron star mergers and how to study them*, [Living Reviews in Relativity **23** \(2020\) 4 \[1909.06085\]](#).
- [33] C.D. Bochenek, V. Ravi, K.V. Belov, G. Hallinan, J. Kocz, S.R. Kulkarni et al., *A fast radio burst associated with a Galactic magnetar*, [Nature **587** \(2020\) 59 \[2005.10828\]](#).
- [34] R. Margutti, B.D. Metzger, R. Chornock, I. Vurm, N. Roth, B.W. Grefenstette et al., *An Embedded X-Ray Source Shines through the Aspherical AT 2018cow: Revealing the Inner Workings of the Most Luminous Fast-evolving Optical Transients*, [ApJ **872** \(2019\) 18 \[1810.10720\]](#).
- [35] K.E. O’Donnell and T.R. Slatyer, *Constraints on dark matter with future MeV gamma-ray telescopes*, [Phys. Rev. D **111** \(2025\) 083037 \[2411.00087\]](#).
- [36] M. Cirelli and A. Kar, *Prospects of future MeV telescopes in probing weak-scale Dark Matter*, [arXiv e-prints \(2025\) arXiv:2503.04907 \[2503.04907\]](#).
- [37] P. De la Torre Luque, S. Balaji and J. Silk, *New 511 keV Line Data Provide Strongest sub-GeV Dark Matter Constraints*, [ApJ **973** \(2024\) L6 \[2312.04907\]](#).
- [38] P. De la Torre Luque, S. Balaji, M. Fairbairn, F. Sala and J. Silk, *511 keV Galactic Photons from a Dark Matter Spike*, [arXiv e-prints \(2024\) arXiv:2410.16379 \[2410.16379\]](#).
- [39] R.D. Peccei and H.R. Quinn, *CP conservation in the presence of pseudoparticles*, [Phys. Rev. Lett. **38** \(1977\) 1440](#).
- [40] W.C. Haxton, X. Liu, A. McCutcheon and A. Ray, *A Continuous Galactic Line Source of Axions: The Remarkable Case of ^{23}Na* , [arXiv e-prints \(2025\) arXiv:2505.03038 \[2505.03038\]](#).
- [41] J. Jaeckel and A. Ringwald, *The Low-Energy Frontier of Particle Physics*, [Annual Review of Nuclear and Particle Science **60** \(2010\) 405 \[1002.0329\]](#).
- [42] A. Lella, E. Ravensburg, P. Carena and M.C.D. Marsh, *Supernova limits on QCD axionlike particles*, [Phys. Rev. D **110** \(2024\) 043019 \[2405.00153\]](#).
- [43] S. Xiao, S.-L. Xiong, Y. Wang, S.-N. Zhang, H. Gao, Z. Zhang et al., *A Robust Estimation of Lorentz Invariance Violation and Intrinsic Spectral Lag of Short Gamma-Ray Bursts*, [ApJ **924** \(2022\) L29](#).

- [44] C. Keith, D. Hooper, T. Linden and R. Liu, *Sensitivity of future gamma-ray telescopes to primordial black holes*, *Phys. Rev. D* **106** (2022) 043003 [2204.05337].
- [45] Z. Xie, B. Liu, J. Liu, Y.-F. Cai and R. Yang, *Limits on the primordial black holes dark matter with future MeV detectors*, *Phys. Rev. D* **109** (2024) 043020 [2401.06440].
- [46] V. Tatischeff, P. Ubertini, T. Mizuno and L. Natalucci, *Orbits and Background of Gamma-Ray Space Instruments*, in *Handbook of X-ray and Gamma-ray Astrophysics*, C. Bambi and A. Sanganello, eds., p. 53 (2022), DOI.
- [47] N. Striebig, R. Leys, I. Peric, R. Caputo, A.L. Steinhebel, Y. Suda et al., *AstroPix4 — a novel HV-CMOS sensor developed for space based experiments*, *Journal of Instrumentation* **19** (2024) C04010.
- [48] R. Caputo, M. Ajello, C.A. Kierans, J.S. Perkins, J.L. Racusin, L. Baldini et al., *All-sky Medium Energy Gamma-ray Observatory eXplorer mission concept*, *Journal of Astronomical Telescopes, Instruments, and Systems* **8** (2022) 044003 [2208.04990].

newASTROGAM collaboration author list as submitted to ESA's M8 call (national contact points in bold)

Lead proposer ESA M8 step 1 proposal: David Berge

Proposal co-PIs: David Berge, M. Nicola Mazziotta, Uwe Oberlack, Vincent Tatischeff, Marco Tavani

M. Ackermann¹, M. Ajello², R. Aloisio³, E. Amato⁴, G. Ambrosi⁵, A. Argan⁶, P. von Ballmoos⁷, F. C. T. Barbato³, M. Barschke¹, R. Battiston⁸, **D. Berge**^{1,9}, N. Berger¹⁰, E. Bissaldi¹¹, M. Branchesi³, S. Brandt¹², T. J. Brandt¹³, C. Budtz-Jørgensen¹², A. Bulgarelli¹⁴, S. Buson^{1,15}, F. Capitanio¹⁶, **R. Caputo**¹⁷, M. Cardillo¹⁶, C. Casentini¹⁶, M. Chernyakova¹⁸, S. Ciprini¹⁹, A. Coleiro²⁰, E. Costa¹⁶, **E. Costantini**¹³, **R. Curado da Silva**²¹, S. Cutini⁵, A. De Angelis²², M. Del Santo²³, D. Della Monica Ferreira¹², D. de Martino²⁴, I. De Mitri³, E. de Oña Wilhelmi¹, N. de Séréville²⁵, A. Di Giovanni³, L. Di Venere²⁶, M. Duranti⁵, S. Fegan²⁷, M. Feroci¹⁶, V. Fioretti¹⁴, A. Franckowiak²⁸, L. Foffano¹⁶, S. Funk²⁹, S. Gallego¹⁰, F. Gargano²⁶, D. Gascón³⁰, D. Gasparrini¹⁹, S. Germani³¹, G. Ghirlanda³², I. Gregor¹, I. A. Grenier³³, L. Hanlon³⁴, D. Hartmann², M. Hernanz³⁵, A. Hornstrup¹², **R. Hudec**³⁶, L. Huth¹, A. Ingram³⁷, R. Iuppa⁸, L. Izzo²⁴, **M. Kachelriess**³⁸, L. Kuiper¹³, **I. Kuvvetli**¹², P. Laurent³³, M. Lemoine-Goumard³⁹, O. Limousin³³, M. Linares³⁸, **T. Linden**⁴⁰, **E. Lindfors**⁴¹, J. Lommler¹⁰, F. Longo⁴², F. Loparco⁴³, N. Lund¹², L. Marcotulli¹, A. Marcowith⁴⁴, K. Mannheim¹⁵, **M. N. Mazziotta**²⁶, **S. McBreen**³⁴, A. Meuris³³, M. Michalska⁴⁵, A. Morselli¹⁸, D. Murphy³⁴, **K. Nakazawa**⁴⁶, L. Nava³², F. Nozzoli⁸, U. Oberlack¹⁰, A. Obertelli⁴⁷, F. Oikonomou³⁸, S. Paltani⁴⁸, A. Papitto⁴⁹, N. Parmiggiani¹⁴, M. Pearce⁵⁰, I. Peric⁵¹, G. Piano¹⁶, F. Piron⁴⁴, C. Pittori⁴⁹, M. Pohl^{1,52}, R. Rando²², **J. Rico**⁵³, S. Ringsborg Howalt Owe¹², G. Rodriguez-Fernandez¹⁹, F. Ryde⁵⁰, O. Salafia³², A. Santangelo⁵⁴, T. Sbarrato³², D. Serini²⁶, T. Siegert¹⁵, K. Skup⁴⁵, S. Spannagel¹, G. Tagliaferri³², **V. Tatischeff**²⁵, M. Tavani¹⁶, F. Tavecchio³², A. Tykhonov⁴⁸, A. Ulyanov³⁴, S. Vercellone³², F. Verrecchia⁴⁹, J. Vink⁵⁵, S. Vitek³⁶, V. Vittorini¹⁶, R. Walter⁴⁸, J. Wilms²⁸, R. Woolf⁵⁶, **X. Wu**⁵⁷, **S. Zane**⁵⁸, **A. Zdziarski**⁵⁹, S. Zhu¹, A. Zoglauer⁶⁰ and P. Zuccon⁸.

¹Deutsches Elektronen-Synchrotron DESY Zeuthen, Germany; ²Department of Physics and Astronomy, Clemson University, Clemson, USA; ³Gran Sasso Science Institute and INFN LNGS, Italy; ⁴Istituto Nazionale di Astrofisica (INAF) OAA Arcetri, Italy; ⁵Istituto Nazionale di Fisica Nucleare (INFN) Perugia, Italy; ⁶INAF Headquarters Roma, Italy; ⁷IRAP Toulouse, France; ⁸University and INFN Trento, Italy; ⁹Humboldt University Berlin, Germany; ¹⁰University Mainz, Germany; ¹¹Politecnico and INFN Bari, Italy; ¹²DTU Space, National Space Institute, Technical University of Denmark; ¹³SRON Netherlands Institute for Space Research, Netherlands ¹⁴INAF-OAS Bologna, Italy; ¹⁵University Würzburg, Germany; ¹⁶INAF-IAPS Roma, Italy; ¹⁷NASA Goddard Space Flight Center, MD, USA; ¹⁸Dublin City University, Ireland; ¹⁹INFN Roma Tor Vergata, Italy; ²⁰APC, Université de Paris, CNRS, France; ²¹LIP, Departamento de Física Universidade de Coimbra, Portugal; ²²University and INFN Padova, Italy; ²³INAF-IASF Palermo, Italy; ²⁴INAF-OA Capodimonte, Italy; ²⁵University of Paris-Saclay, CNRS, IJCLab, France; ²⁶INFN Bari, Italy; ²⁷LLR, École Polytechnique, CNRS, Institut Polytechnique de Paris, France; ²⁸University Bochum, Germany; ²⁹University Erlangen, Germany; ³⁰Institute of Cosmos Sciences of the University of Barcelona (ICCUB), Spain; ³¹University and INFN Perugia, Italy; ³²INAF-OAB Merate, Italy; ³³CEA/IRFU, Université Paris-Saclay, France; ³⁴University College Dublin, Ireland; ³⁵ICE-CSIC and IEEC, Barcelona, Spain; ³⁶Czech Technical University Prague, Czech Republic; ³⁷Newcastle University, Great Britain; ³⁸NTNU, Trondheim, Norway; ³⁹Univ. Bordeaux, CNRS, LP2i Bordeaux,

France; ⁴⁰Physics Department, Stockholm University, Sweden; ⁴¹Department of Physics and Astronomy, University of Turku, Finland; ⁴²University and INFN Trieste, Italy; ⁴³University and INFN Bari, Italy; ⁴⁴LUPM, Université Montpellier, CNRS, France; ⁴⁵Space Research Center, Polish Academy of Sciences, Warszawa, Poland; ⁴⁶Space Astronomy Lab, Nagoya University, Japan; ⁴⁷University Darmstadt, Germany; ⁴⁸Department of Astronomy, University of Geneva, Switzerland; ⁴⁹INAF-OAR Roma, Italy; ⁵⁰Royal Institute of Technology (KTH), Stockholm, Sweden; ⁵¹Karlsruhe Institute for Technology, Germany; ⁵²University Potsdam, Germany; ⁵³Institut de Física d'Altes Energies (IFAE), Barcelona, Spain; ⁵⁴IAAT, University Tübingen, Germany; ⁵⁵University of Amsterdam, Netherlands; ⁵⁶Naval Research Laboratory Washington DC, USA; ⁵⁷Department of Nuclear and Particle Physics, University of Geneva, Switzerland; ⁵⁸Mullard Space Science Laboratory, University College, London, UK; ⁵⁹Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, Warszawa, Poland; ⁶⁰University of California at Berkeley, Space Sciences Laboratory, USA.